

Wigner Functions in Curved Space–Time and Quantum Corrections to Thermal Equilibrium

Oleg A. Fonarev*

Racah Institute of Physics, The Hebrew University

Jerusalem 91904, Israel

E-mail: OLG@vms.huji.ac.il

The Wigner function is known to be a very useful tool in exploring the semiclassical limit of the quantum theory. It is defined as a Fourier representation of the density matrix ρ with respect to the difference of coordinates¹:

$$f_w(q, p) = (\pi\hbar)^{-1} \int_{-\infty}^{\infty} dQ e^{-2iQp/\hbar} \langle q + Q | \rho | q - Q \rangle . \quad (1)$$

At a high temperature or after coarse graining, the Wigner function gives us directly quantum corrections to a distribution function in the classical phase space (the Boltzmann distribution if the system is in thermal equilibrium).

In curved space–time the notion of the Fourier transformation becomes ambiguous because it is not invariant under general coordinate transformations. Different approaches to solve this problem have been proposed^{2–5}. I will follow the approach by [5,6] which uses the formalism of the tangent bundles. Consider any point x on a space–time manifold $\mathcal{M}$. Let $\varphi(x)$ be a (real) scalar field on the manifold. Let us introduce the tangent space $\mathcal{T}_x(\mathcal{M})$ at point x and define field $\Phi(x, y)$ on the tangent bundle as follows:

$$\Phi(x, y) = \exp(y^\alpha \hat{\nabla}_\alpha) \varphi(x) . \quad (2)$$

Here the operator $\hat{\nabla}_\alpha = \nabla_\alpha - \Gamma_{\alpha\gamma}^\beta y^\gamma \frac{\partial}{\partial y^\beta}$ is the horizontal lift of the derivative operator to the tangent bundle⁷. The generalized Wigner function is now defined as follows:

$$f(x, p) = (\pi\hbar)^{-4} \sqrt{-g(x)} \int_{\mathcal{T}_x(\mathcal{M})} d^4 y e^{-2iy^\alpha p_\alpha/\hbar} \langle \Phi(x, -y) \Phi^\dagger(x, y) \rangle . \quad (3)$$

Here the brackets mean the averaging with a density matrix on an initial Cauchy hypersurface. In the Minkowski space–time, the tangent space $\mathcal{T}_x(\mathcal{M})$ coincides with the base space $\mathcal{M}$, the exponent in Eq.(2) becomes the shift operator, i.e. $\Phi(x, y) = \varphi(x + y)$ and we get the usual definition of the relativistic Wigner function⁸. In curved space–time, the definition (3) is explicitly covariant and allows us to easily derive equations describing the evolution of the generalized Wigner function in the phase space, given a field equation for the field φ .

Suppose that φ obeys the covariant Klein–Gordon equation with the conformal coupling to gravity:

$$\left(\nabla^\alpha \nabla_\alpha + m^2/\hbar^2 - \frac{1}{6} R \right) \varphi = 0 , \quad (4)$$

where R is the scalar curvature. We then get two equations for the Wigner function^{2–6}:

$$\begin{aligned} (m^2 - p^\alpha p_\alpha) f &= -\frac{\hbar^2}{4} D^\alpha D_\alpha f - \\ &- \hbar^2 \left(\frac{1}{6} R + \frac{1}{12} R_{\alpha\mu\beta\nu} p^\mu p^\nu \frac{\partial^2}{\partial p_\alpha \partial p_\beta} + \frac{1}{4} R_{\mu\nu} p^\mu \frac{\partial}{\partial p_\nu} \right) f + O(\hbar^4) , \end{aligned} \quad (5)$$

*talk given at the 3rd International Wigner Symposium, 5th–11th September 1993, Oxford, UK.

$$\begin{aligned}
p^\alpha D_\alpha f = & \hbar^2 \left(\frac{1}{6} R_{\nu\beta\mu\alpha} p^\mu \frac{\partial^2}{\partial p_\alpha \partial p_\beta} D^\nu - \frac{1}{24} R_{\alpha\mu\beta\nu;\sigma} p^\mu p^\nu \frac{\partial^3}{\partial p_\alpha \partial p_\beta \partial p_\sigma} + \right. \\
& \left. + \frac{1}{12} R_{\alpha}^{\nu} \frac{\partial}{\partial p_\alpha} D_\nu - \frac{1}{24} R_{\alpha\beta;\nu} p^\nu \frac{\partial^2}{\partial p_\alpha \partial p_\beta} - \frac{1}{24} R_{;\alpha} \frac{\partial}{\partial p_\alpha} \right) f + O(\hbar^4), \tag{6}
\end{aligned}$$

where $D_\alpha = \nabla_\alpha + \Gamma_{\alpha\beta}^\gamma p_\gamma \frac{\partial}{\partial p_\beta}$ is the horizontal lift of the derivative operator to the *cotangent* bundle⁷.

Eq.(5) is a generalized mass-shell constraint, and it implies that in the semiclassical limit, the structure of the Wigner function must be as follows³:

$$f = F_0 \delta(m^2 - p^2) + \hbar^2 F_1 \delta'(m^2 - p^2) + \hbar^2 F_2 \delta''(m^2 - p^2) + O(\hbar^4). \tag{7}$$

All the F_n 's in (7) are easily found from eq.(5) and they are expressed^{3,6} in terms of certain local differential operators acting to the function F_0 . The function F_0 satisfies the quantum corrected Vlasov equation (6) on the mass-shell. F_0 is decomposed to classical and quantum parts:

$$F_0 = F_{cl} + \hbar^2 F_{qu} + O(\hbar^4), \tag{8}$$

where F_{qu} is found by integrating the classical distribution function F_{cl} along classical trajectories and is in general nonlocal in the phase space.

Thus, given a classical distribution function $F_{cl}(x, p)$ as a solution to the Vlasov equation, one is able, in principle, to evaluate the semiclassical expansion of the generalized Wigner function and of any physical observable expressible in terms of the Wigner function. For example, the number-flux $N_\alpha(x)$ and the stress-energy vector $T_{\alpha\beta}(x)$ are given by^{2,6}:

$$N_\alpha(x) = \int \frac{d^4 p}{\sqrt{-g(x)}} p_\alpha f(x, p), \tag{9}$$

$$\begin{aligned}
T_{\alpha\beta}(x) = & \int \frac{d^4 p}{\sqrt{-g(x)}} p_\alpha p_\beta f(x, p) + \\
& + \hbar^2 \left(\frac{1}{6} R_{\alpha\beta}(x) + \frac{1}{12} (\nabla_\alpha \nabla_\beta - g_{\alpha\beta}(x) \nabla^\nu \nabla_\nu) \right) \int \frac{d^4 p}{\sqrt{-g(x)}} f(x, p). \tag{10}
\end{aligned}$$

Let us now consider a stationary space-time admitting a global time-like Killing vector ξ^α . In that case, any function, $F_{cl}(E)$, of the constant of motion, $E = \xi^\alpha p_\alpha$, is a solution of the Vlasov equation. The classical stress-energy tensor calculated with such distribution has a perfect fluid's structure. The quantum corrections of the lowest adiabatic order to this distribution were found in Ref.[9] and they possess the following properties: a) all the quantum corrections are local in the phase space and they are obtained by acting a certain differential operator to F_{cl} . This means that physical observables are expressed in terms of momenta of the classical distribution function; b) the stress-energy tensor has the following structure:

$$T_{\alpha\beta} = (\varepsilon + p) v_\alpha v_\beta - p g_{\alpha\beta} - \Pi_{\alpha\beta}, \tag{11}$$

which is not longer a perfect fluid's one : the anisotropic pressure tensor $\Pi_{\alpha\beta}$ appears ($\Pi_\alpha^\alpha = v^\alpha \Pi_{\alpha\beta} = 0$); c) the eigenvector of the stress-energy tensor, v_α , doesn't coincide with the hydrodynamical velocity $u_\alpha = N_\alpha (N^\nu N_\nu)^{-1/2}$. This leads to the nonvanishing heat flow: $I_\alpha = (\varepsilon + p)(v_\alpha - u_\alpha)$. I_α is proportional to the combination $N_0/6N_2 - N_1/4N_3$, where $N_k = \int_0^\infty dE E^k F_{cl}(E)$. For the Boltzmann or Bose-Einstein distributions this combination doesn't vanish.

Only in static space-times (i.e. for nonrotating systems) the quantum corrections preserve

the perfect fluid's structure with the vanishing heat flow.

Let us next consider a more general case of conformally stationary (or conformally static) space-times. The definition (3) of the generalized Wigner function given allows one to easily link Wigner functions in two conformally related manifolds. Under the conformal transformation, $g_{\alpha\beta}(x) \mapsto a(x)^2 g_{\alpha\beta}(x)$, $\varphi(x) \mapsto \varphi(x)/a(x)$, the Wigner function transforms as follows¹⁰:

$$f \mapsto a^2(1 + \hbar^2 \hat{A} + O(\hbar^4))f, \quad (12)$$

where $\hat{A}$ is a certain differential operator involving derivatives of the conformal factor $a(x)$. If one knows a Wigner function in a stationary space-time, one is able to compute a Wigner function in a conformally stationary space-time. We have found¹⁰ by this method the quantum corrections to a class of distribution functions isotropic in the momentum space¹¹, $F_{cl} = F(u^\alpha p_\alpha, x)$.

The Robertson-Walker Universes are examples of conformally static (and also conformally flat) space-times. The metric tensor takes the form: $g_{\alpha\beta} = a(t)^2 \gamma_{\alpha\beta}$ where $\gamma_{\alpha\beta}$ is the metric tensor of a spherically symmetrical static space-time, and t is the conformal time. The classical distribution function compatible with the symmetry of the space-time is any function of the three-momentum¹¹. The quantum corrections to such distributions were first found by [12] through a different method. The structure of the quantum-corrected stress-energy tensor is a perfect fluid's one in this case. We also found¹⁰ that the quantum corrected energy density takes the following form:

$$\varepsilon(t) = \varepsilon_{cl}(t) + \frac{1}{\kappa} \frac{m^2}{M_{pl}^2} \mathcal{K}\left(\frac{m}{\Theta(t)}\right) G_0^0(t) + O(\hbar^4) \quad (13)$$

where $\mathcal{K}$ is a certain function of the ratio of the particles mass m to the local temperature $\Theta(t) = T/a(t)$, M_{pl} is the Planck mass and G_0^0 is the zero-zero component of the Einstein tensor. If one substitutes Expr. (13) into the Einstein equation, one finds that the effective gravitational "constant" appears which varies with time:

$$\frac{1}{\kappa_{eff}(t)} = \frac{1}{\kappa} \left(1 - \frac{m^2}{M_{pl}^2} \mathcal{K}\left(\frac{m}{\Theta(t)}\right)\right). \quad (14)$$

The variation is negligible at present but it could lead to important physical effects in the early Universe.

¹ E. P. Wigner, Phys. Rev. **40**, 749 (1932).

² J. Winter, Phys. Rev. D **32**, 1871 (1985).

³ E. Calzetta, S. Habib, and B. L. Hu, Phys. Rev. D **37**, 2901 (1988).

⁴ H. E. Kandrup, Phys. Rev. D **37**, 2165 (1988).

⁵ O. A. Fonarev, Izv. VUZov. Fizika (USSR) **9**, 47 (1990).

⁶ O. A. Fonarev, preprint gr-qc/9309005; to appear in J.Math.Phys. (USA).

⁷ K. Yano and Sh. Ishihara, *Tangent and Cotangent Bundles: Differential Geometry* (Pure and Applied Mathematics: A Series of Monographs and Textbooks) (Marcel Dekker, Inc. , New York, 1973).

⁸ S. R. de Groot, W. A. van Leeuwen, and Ch. G. van Weert, *Relativistic Kinetic Theory* (North-Holland, Amsterdam, 1980).

⁹ O. A. Fonarev, Phys. Lett. A **152**, 153 (1991).

- ¹⁰ O. A. Fonarev, in preparation.
- ¹¹ J. Ehlers, P. Geren and R. K. Sachs, J. Math. Phys. **9**, 1344 (1968).
- ¹² K. Pirk and G. Börner, Class. Quantum Grav. **6**, 1855 (1989).